



**FP-10: Increase the proportion of sexually active persons aged 15 to 19 years who use condoms to both effectively prevent pregnancy and provide barrier protection against disease.**

FP-10.1 & 10.2 Increase the proportion of sexually active teens aged 15 to 19 years who use a condom at first intercourse.

10.1 females	Target: 74.8%	Baseline: 2006–10, 68.0%
10.2 males	Target: 87.6%	Baseline: 2006–10, 79.6%

FP-10.3 & 10.4 Increase the proportion of sexually active teens aged 15 to 19 years who use a condom at last intercourse.

10.3 females	Target: 55.6%	Baseline: 2006–10, 50.5%
10.4 males	Target: 81.5%	Baseline: 2006–10, 74.1%

Data source: National Survey of Family Growth (NSFG), CDC, NCHS.

**FP-11: Increase the proportion of sexually active persons aged 15 to 19 years who use condoms and hormonal or intrauterine contraception to both effectively prevent pregnancy and provide barrier protection against disease.**

FP-11.1 & 11.2 Increase the proportion of sexually active teens aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse.

11.1 females	Target: 15.4 %	Baseline: 2006–10, 14.0%
11.2 males	Target: 17.3 %	Baseline: 2006–10, 15.7%.

FP-11.3 & 11.4 Increase the proportion of sexually active teens aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse.

11.3 females	Target: 20.1 %	Baseline: 2006–10, 18.3%
11.4 males	Target: 35.3 %	Baseline: 2006–10, 32.1%

Data source: National Survey of Family Growth (NSFG), CDC, NCHS.

**FP-12: Increase the proportion of adolescents who received formal instruction on reproductive health topics before they were 18 years old.**

FP-12.1 Abstinence—Females	Target: 97.6 %	Baseline: 2006–10, 88.7%
FP-12.2 Abstinence—Males	Target: 90.8 %	Baseline: 2006–10, 82.5%
FP-12.3 Birth control methods—Females	Target: 77.6 %	Baseline: 2006–10, 70.5%
FP-12.4 Birth control methods—Males	Target: 66.9 %	Baseline: 2006–10, 60.8%
FP-12.5 HIV/AIDS prevention—Females	Target: 98.2 %	Baseline: 2006–10, 89.3%
FP-12.6 HIV/AIDS prevention—Males	Target: 96.7 %	Baseline: 2006–10, 87.9%
FP-12.7 STDs—Females	Target: 95.8 %	Baseline: 2006–10, 93.8%
FP-12.8 STDs—Males	Target: 93.8 %	Baseline: 2006–10, 91.8%

Data source: National Survey of Family Growth (NSFG), CDC, NCHS.

**FP-13: Increase the proportion of adolescents who talked to a parent or guardian about reproductive health topics before they were 18 years old.**

FP-13.1 Abstinence—Females	Target: 67.5 %	Baseline: 2006–10, 61.4%
FP-13.2 Abstinence—Males	Target: 45.3 %	Baseline: 2006–10, 41.2%
FP-13.3 Birth control methods—Females	Target: 56.1 %	Baseline: 2006–10, 51.0%
FP-13.4 Birth control methods—Males	Target: 32.1 %	Baseline: 2006–10, 29.2%
FP-13.5 HIV/AIDS prevention—Females	Target: 45.0 %	Baseline: 2006–10, 40.9%
FP-13.6 HIV/AIDS prevention—Males	Target: 41.6 %	Baseline: 2006–10, 37.8%
FP-13.7 STDs—Females	Target: 59.6 %.	Baseline: 2006–10, 54.2%
FP-13.8 STDs—Males	Target: 52.9 %	Baseline: 2006–10, 48.1%

Data source: National Survey of Family Growth (NSFG), CDC, NCHS.

**Topic Area: Sexually Transmitted Diseases****STD-5: Reduce the proportion of females aged 15 to 44 years who have ever required treatment for pelvic inflammatory disease (PID).**

Target: 3.8%

Baseline: 2006-10, 4.2%

Data source: National Survey of Family Growth (NSFG), CDC, NCHS

**Topic Area: HIV****HIV-14: Increase the proportion of adolescents and adults who have been tested for HIV in the past 12 months.**

HIV-14.1 Adolescents and adults.

Target: 73.6%

Baseline: 2006-10, 66.9%

HIV-14.2 (Developmental) Men who have sex with men (MSM).

Target: 68.4%

Baseline: 2006-08, 62.2%

HIV-14.3 Pregnant women.

Target: 79.2%

Baseline: 2006-10, 72.0%

**Topic Area: Maternal, Infant, and Child Health****MICH-17: Reduce the proportion of persons aged 18 to 44 years who have impaired fecundity (i.e., a physical barrier preventing pregnancy or carrying a pregnancy to term).**

MICH-17.1 Reduce the proportion of women aged 18 to 44 years who have impaired fecundity.

Target: 11.4%

Baseline: 2002, 12.7%

MICH-17.2 (Developmental) Reduce the proportion of men aged 18 to 44 years who have impaired fecundity.

Data source: National Survey of Family Growth (NSFG), CDC, NCHS.